

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 0 945 066 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
29.09.1999 Bulletin 1999/39

(51) Int Cl.⁶: **A01N 65/00**
// (A01N65/00, 65:00, 63:00,
27:00)

(21) Application number: **99302286.2**

(22) Date of filing: **25.03.1999**

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: **26.03.1998 US 79505 P**
23.03.1999 US

(71) Applicant: **Safergro Laboratories, Inc.**
Ventura, CA 93003 (US)

(72) Inventors:
• **Hsu, Hsinhung John**
Ventura, CA 93003 (US)
• **Chang, Chun-Hua Lily**
Ventura, CA 93003 (US)
• **Zhou, Jian NMI**
Ventura, CA 93003 (US)

(74) Representative: **Kirkham, Nicholas Andrew et al**
Graham Watt & Co.,
Riverhead
Sevenoaks, Kent TN13 2BN (GB)

(54) **Natural pesticide containing garlic**

(57) The invention describes a synergistic effect when garlic oil or extract is combined with essential oils which results in an improved insecticide/fungicide which is natural and contains no chemical additives. Essential oils are defined in this application to be volatile liquids obtained from plants and seeds including cotton seed

oil, soybean oil, cinnamon oil, corn oil, cedar oil, castor oil, clove oil, geranium oil, lemongrass oil, linseed oil, mint oil, sesame oil, thyme oil, rosemary oil, anise oil, basil oil, camphor oil, citronella oil, eucalyptus oil, fennel oil, ginger oil, grapefruit oil, lemon oil, mandarin oil, orange oil, pine needle oil, pepper oil, rose oil, tangerine oil, tea tree oil, tee seed oil, mineral oil and fish oil.

EP 0 945 066 A1

Description

[0001] This invention relates to a composition of matter used as a natural pesticide and a method for inhibiting the growth of bacteria, fungi and insect pests.

[0002] Garlic (*Allium sativum* Linn.) and/or its extract have been reported to have antibacterial and/or antifungal properties. It is known that Allicin isolated from the cloves of garlic had antibacterial properties against both Gram positive and Gram negative bacteria. Further, aqueous extracts of garlic have been reported to inhibit the growth of a variety of yeast-like fungi in the genera *Candida*, *Cryptococcus*, *Rhizotoruto*, *Torulopsis* and *Trichosporon*. It has also been previously reported that garlic extract and chips inhibit the growth of fungi such as *Candida albicans*, *Aspergillus fumigatus* and *Aspergillus parasiticus*. Because of its antifungal and antibacterial properties, garlic or its extract have been used as pesticides to control plant diseases such as mildew. It has also been used as an insecticide to control plant insects such as army worms, aphids and Colorado beetles. Most recently, a method used to repel mosquitos using garlic extract and water was granted US Pat. No. 5,733,552 issued to Anderson et. al.

[0003] The present invention is directed to a composition of matter which comprises garlic extract and essential oils. This combination of natural ingredients, when applied to plants, has superior anti-fungal and anti-bacterial qualities, than if applied separately.

[0004] A combination of garlic extract and essential oil has a synergistic effect which significantly increases the effectiveness of garlic and/or garlic extract. The ideal ratio of garlic to essential oil is 10-70% garlic extract to 90-30% essential oil. However, a ratio of 5-98% garlic extract to 95-2% essential oil can be used.

[0005] Garlic extract can be obtained by the blending and mixing of garlic cloves with water, oil or organic solvents. The mixture is then filtered to obtain garlic extract. In addition to essential oils, mineral oil and fish oil can also be used.

Definitions

[0006] Essential Oil is defined as a subtle, volatile liquid obtained from plants and seeds, including but not limited to cotton seed oil, soybean oil, cinnamon oil, corn oil, cedar oil, castor oil, clove oil, geranium oil, lemongrass oil, linseed oil, mint oil, sesame oil, thyme oil, rosemary oil, anise oil, basil oil, camphor oil, citronella oil, eucalyptus oil, fennel oil, ginger oil, grapefruit oil, lemon oil, mandarin oil, orange oil, pine needle oil, pepper oil, rose oil, tangerine oil, tea tree oil and tea seed oil.

[0007] Garlic extract is defined as any liquid removed from cloves of garlic and may therefore include garlic oil and water. Garlic extract has the same meaning as garlic juice.

[0008] Disease index represents the severity of the disease present on a plant. This is a subjective assessment and is categorized in the following manner:

Scale	% Infection
0	0
1	1-10
2	11-20
3	21-50
4	51-100

[0009] The disease index is calculated by the following formula:

$$DI = \frac{[(S_0L_0) + (S_1L_1) + (S_2L_2) + (S_3L_3) + (S_4L_4)]}{L_{total} \times 4} \times 100$$

[0010] Where DI=Disease index

S=Scale category (0, 1, 2, 3 or 4)

L=number of leaves per scale category

[0011] The degree of control represents the efficiency of the product in controlling the disease and is calculated by the following formula:

$$DC = \frac{DI_{\text{non-treated area}} - DI_{\text{treated area}}}{DI_{\text{non-treated area}}} \times 100$$

5 **[0012]** A degree of control of 0 represents no control and 100 represents total control of the disease.

Preparation of Garlic Extract

10 **[0013]** Garlic extract was prepared by thoroughly mixing two thousand grams of garlic cloves with 4,000 grams of water in a blender. The mixture was then filtered with the filtrate collected as garlic extract.

Samples 1-4

15 **[0014]** Samples 1-4 were mixed as indicated in the following chart at ambient conditions:

Composition of Samples 1-4 (all values in grams)				
	Sample 1	Sample 2	Sample 3	Sample 4
Garlic Extract	600	850	600	600
Cotton Seed Oil	300	--	250	--
Cinnamon Oil	--	50	50	--
Mineral Oil	--	--	--	300
Sodium Lauryl Sulfate	100	100	100	100

20 **[0015]** Sodium Lauryl Sulfate is used to emulsify the garlic extract with either an essential oil or mineral oil.

EXPERIMENTS

30

Experiment A - Powdery Mildew Disease on Cucumber Plants

35 **[0016]** Cucumber plants were grown for 14 days in a greenhouse. The leaves were then inoculated with a spore suspension of powdery mildew disease. Three weeks after the spore inoculation, all cucumber plants developed powdery mildew disease. Fifteen cucumber plants were selected for testing. The plants were randomly assigned into 5 groups having 3 plants per group. Five spray compositions were prepared. Each of the following, Sample 1, Garlic Extract, Cotton Seed Oil and Sodium Lauryl Sulfate were diluted one part to 49 parts water. The fifth spray composition was water.

40 **[0017]** Each group of plants then received one of the above-mentioned compositions by foliar spray treatment twice with a 7 day interval between each treatment. The spray treatments consisted of totally wetting the plant leaves.

[0018] Seven days after the second application, the plants were examined for powdery mildew disease. The disease indices were recorded to show the severity of the disease and are reported below:

Test Results 7 Days after Second Spray Application		
	Disease Index	Degree of Control
Sample 1	1.3	97.2%
Garlic Extract	5.0	89.4%
Cotton Seed Oil	8.0	83.1%
Sodium Lauryl Sulfate	31.8	32.6%
Water	47.2	0.0%

55 **[0019]** The results indicate that garlic extract in combination with cotton seed oil (sample 1) is better than garlic extract or cotton seed oil alone on controlling powdery mildew disease on cucumber plants.

Experiment B - Powdery Mildew Disease on Cucumber Plants

[0020] The testing in Experiment A was repeated with 28 cucumber plants divided into 7 groups (Groups 6-12), each group having 4 plants. Seven spray compositions were prepared. Each of the following: Sample 1, Sample 2, Garlic Extract, Cotton Seed Oil, Cinnamon Oil and Sodium Lauryl Sulfate, were diluted one part to 99 parts water. The seventh spray composition was water. The seven groups received the foliar spray treatments twice with a 7 day interval between each treatment.

[0021] Three days after each application, the plants were examined for powdery mildew disease. The disease indices were recorded to show the severity of the disease and are reported as follows:

Test Results 3 Days after Application				
	1 st Application		2 nd Application	
	Disease Index	Degree of Control	Disease Index	Degree of Control
Sample 1	14.6	77.4%	10.0	81.3%
Sample 2	12.5	80.6%	6.7	87.5%
Garlic Extract	27.1	58.2%	20.0	62.5%
Cotton Seed Oil	56.3	12.9%	43.3	18.8%
Cinnamon Oil	56.3	12.9%	50.0	6.2%
Sodium Lauryl Sulfate	60.4	6.4%	51.7	3.1%
Water	64.4	0.0%	53.3	0.0%

[0022] The results indicate that the combination of garlic extract and either cottonseed oil (sample 1) or cinnamon oil (sample 2) inhibit the growth of powdery mildew disease more effectively than applications not in combination.

Experiment C - Powdery Mildew Disease on Cucumber Plants

[0023] The testing in Experiment B was repeated; this time with each foliar spray application diluted with 199 parts water instead of 99 parts water.

[0024] Three days after each application, the plants were examined for powdery mildew disease. The disease indices were recorded to show the severity of the disease and are reported as follows:

Test Results 3 Days after Application				
	1 st Application		2 nd Application	
	Disease Index	Degree of Control	Disease Index	Degree of Control
Sample 1	25.0	61.3%	20.0	62.5%
Sample 2	25.0	61.3%	25.0	53.1%
Garlic Extract	33.3	48.6%	26.7	50.0%
Cotton Seed Oil	60.4	6.5%	45.0	15.6%
Cinnamon Oil	56.3	12.9%	51.7	3.1%
Sodium Lauryl Sulfate	62.5	3.2%	51.7	3.1%
Water	64.6	0.0%	53.3	0.0%

[0025] The results indicate that the combination of garlic extract and either cottonseed oil (sample 1) or cinnamon oil (sample 2) inhibit the growth of powdery mildew disease more effectively than either garlic extract, cottonseed oil or cinnamon oil separately. The increased dilution of each spray composition with water did reduce the effectiveness of controlling powdery mildew disease.

Experiment D - Powdery Mildew Disease on Red Rose Plants

[0026] Red rose plants were grown in a greenhouse having powdery mildew disease. In order to test the effectiveness of the combination of garlic extract and cottonseed oil (sample 1) on controlling the disease, the plants were sprayed once with a solution consisting of sample 1 diluted with 50 times water. The control plants were sprayed with water only. The plants were investigated 10 days after the spray application.

The Effectiveness of Sample 1 on Controlling Powdery Mildew

[0027]

Disease on Red Rose Plants		
	Disease Index	Degree of Control
Sample 1	15	66.7%
Garlic Extract	20	55.6%
Cotton Seed Oil	25	44.4%
Water Only	45	0.0%

[0028] Sample 1, a combination of garlic extract and cotton seed oil diluted with 50 times water, was more effective on controlling powdery mildew disease on red rose plants than garlic extract or cotton seed oil individually.

Experiment E - Powdery Mildew Disease on Yellow Rose Plants

[0029] Example D was repeated using yellow rose plants for test and the plants were sprayed with sample 1 diluted with 50 times water, three times at 1 week intervals. The control plants were sprayed with water only. The plants were investigated 10 days after each spray application.

Test Results 10 Days after Application for Controlling Powdery Mildew Disease on Yellow Rose Plants						
	1 st Application		2 nd Application		3 rd Application	
	Disease Index	Degree of Control	Disease Index	Degree of Control	Disease Index	Degree of Control
Sample 1	70	30%	55	45%	15	75%
Garlic extract	80	20%	70	30%	40	60%
Cottonseed oil	85	15%	75	25%	45	55%
Water Only	100	0%	100	0%	100	0%

[0030] Sample 1, a combination of garlic extract and cotton seed oil, when diluted with 50 times water, was more effective on controlling powdery mildew disease on yellow rose plants than garlic extract or cottonseed oil individually.

Experiment F - Powdery Mildew Disease on White Rose Plants

[0031] Example D was repeated using white rose plants for test and the plants were sprayed with sample 1 diluted with 50 times water, twice with a 1 week interval. The control plants were sprayed with water only. The plants were investigated 10 days after both spray applications.

Test Results 10 Days after Application for Controlling Powdery Mildew Disease on White Rose Plants				
	1 st Application		2 nd Application	
	Disease Index	Degree of Control	Disease Index	Degree of Control
Sample 1	55	38.9%	5	87.5%

(continued)

Test Results 10 Days after Application for Controlling Powdery Mildew Disease on White Rose Plants				
	1 st Application		2 nd Application	
	Disease Index	Degree of Control	Disease Index	Degree of Control
Garlic Extract	65	27.8%	20	50.0%
Vegetable Oil	75	16.7%	25	37.5%
Water Only	90	0.0%	40	0.0%

[0032] Sample 1, a combination of garlic extract and cotton seed oil, when diluted with 50 times water, was more effective on controlling powdery mildew disease on white rose plants than either garlic extract or vegetable oil individually.

Experiment G -Powdery Mildew Disease on Cucumber Plants

[0033] Eight cucumber plants with powdery mildew disease were randomly divided into 2 groups having 4 plants each. One group was sprayed with Sample 1 diluted with 49 parts water. The other group of cucumber plants is designated the control group and is only sprayed with water. Both groups of plants were sprayed twice with an interval of 11 days. The disease indices before and after the treatments and the degrees of control are shown below:

The Effectiveness of Sample 1 on Controlling Powdery Mildew Disease on Cucumber Plants

	Disease Index Before Application	1 st Application				2 nd Application			
		- 3 Days -	- 6 Days -	- 11 Days -	- 21Days -				
	Disease Index	Disease Index	Degree of Control	Disease Index	Degree of Control	Disease Index	Degree of Control	Disease Index	Degree of Control
Sample 1	38.8	0	100%	0	100%	17.5	70%	18.8	74%
Water Only	35.0	35.0	0%	46.3	0%	57.5	0%	73.2	0%

Experiment H - Control of White Fly on Cucumber Plants

[0034] Sample 1, diluted with 50 times water, was tested for the control of white fly on cucumber plants. The leaves of cucumber plants were sprayed while adult white flies present, and the mortality rate was tested as follows:

Effectiveness of Sample 2 on Controlling White Fly			
Test No.	White Flies (before spraying)	White Flies Dead (after spraying)	Mortality (%)
1	124	84	67.7
2	74	51	68.9
3	113	62	54.8
4	187	117	62.6

Experiment I - Control of Red Spider Mites on Bush Bean Plants

[0035] Samples 1, 2, 3, and 4 were used to test the control of red spider mites on bush bean plants. The samples, as well as solutions of garlic extract, cotton seed oil, cinnamon oil, and mineral oil were diluted with 100 times water

and sprayed on to bush bean plants which had red spider mites grown on the leaves. The mortality of red spider mites was determined as a percentage in two tests with the results as follows:

Effectiveness Comparison on Red Spider Mites		
Solution (diluted w/100 times water)	Test 1 Mortality (%)	Test 2 Mortality (%)
Sample 1	84	86
Sample 2	88	93
Sample 3	81	84
Sample 4	68	72
Garlic Extract	53	58
Cotton Seed Oil	49	61
Cinnamon Oil	59	64
Mineral Oil	15	28
Water	2	8

Samples 1, 2, 3, and 4 which contain a combination of garlic extract and either an essential oil or mineral oil, were more effective in eradication of red spider mites than using garlic extract, essential oil, or mineral oil separately.

Experiment J - Control of Two Spotted Spider Mites on Bush Bean Plants

[0036] Samples 1, 2, 3, and 4 were used to test the control of two spotted spider mites on bush bean plants. The samples as in Experiment I above, were diluted with 100 times water and sprayed on to bush bean plants which had two spotted spider mites grown on the leaves. The mortality of two spotted spider mites was determined as a percentage in two tests with the results as follows:

Effectiveness Comparison on Two Spotted Spider Mites		
Solution (diluted w/100 times water)	Test 1 Mortality (%)	Test 2 Mortality (%)
Sample 1	78	82
Sample 2	84	91
Sample 3	76	89
Sample 4	65	71
Garlic Extract	53	48
Cotton Seed Oil	36	42
Cinnamon Oil	56	61
Mineral Oil	13	21
Water	5	7

[0037] Samples 1, 2, 3, and 4 which contain a combination of garlic extract and either an essential oil or mineral oil, were more effective in eradication of two spotted spider mites than using garlic extract, essential oil, or mineral oil separately.

[0038] The following experiments relate to suppressing fungal growth (Experiment K) and suppressing fungal germination (Experiment L).

Experiment K - Suppression of *Rhizoctonia* sp., *Trichoderma* sp., and *Botrytis* sp.

[0039] Sample 1 was laboratory tested for the suppression of different fungi. Sample 1 was added to PDA (Potato Dextrose Agar) growth medium solution at the Sample 1:PDA solution ratio between 1:100 and 1:3200. The solutions were then poured into test tubes and inoculated with one of the three different fungi; *Rhizoctonia* sp., *Trichoderma* sp.,

and *Botrytis* sp. The test tubes were then incubated at 25C for 14 days. The growth of the fungi in different treatments were recorded and shown in the following table.

Growth of Three Different Fungi in PDA Solution with Different Concentrations of Sample 1.							
Fungi	0	1/100	1/200	1/400	1/800	1/1600	1/3200
<i>Rhizoctonia</i> sp.	++	+	++	++	++	++	++
<i>Trichoderma</i> sp.	++	-	+	+	+	+	+
<i>Botrytis</i> sp.	++	+	+	+	+	+	+
Legend: ++ Normal Growth + Reduced Growth - No Growth							

[0040] The test results indicate at the Sample 1:PDA ratio of 1:100, Sample 1 suppressed the growth of all fungi tested. However, at weaker concentrations 1:200 - 1:3200, Sample 1 suppressed the growth of both *Trichoderma* sp. and *Botrytis* sp. but not *Rhizoctonia* sp.

Experiment L -Suppression of *Penicillium* sp. *Botrytis* sp. and *Aspergillus* sp. spore germination

[0041] Samples 1, 2, 3, and 4 were tested in the laboratory for the suppression on the germination of spores from 3 different fungi: *Penicillium* sp. *Botrytis* sp. and *Aspergillus* sp. Spore suspensions were prepared and inoculated onto PDA plates. After the inoculation, 1 ml of 1/50 solution of Samples 1, 2, 3, and 4 were added to respective plates receiving the treatment and the plates were horizontally shaken to distribute the solution evenly over the PDA surface. Two sets of control plates were prepared by adding either water or water plus sodium lauryl sulfate solution to the inoculated plates. The plates were then incubated at 26C for 18-20 hours and then examined under a microscope for evidence of spore germination. The number of germinated and ungerminated spores and the percentage of germination are shown in the following table.

5

10

15

20

25

30

35

40

45

50

55

Germination of Spores									
	Penicillium sp.			Botrytis sp.			Aspergillus sp.		
	# Germinated	# un-Germinated	% Germinated	# Germinated	# un-Germinated	% Germinated	# Germinated	# un-Germinated	% Germinated
Sample 1	0	150	0	0	150	0	0	150	0
Sample 2	0	150	0	0	150	0	0	150	0
Sample 3	0	150	0	0	150	0	0	150	0
Sample 4	3	147	2.0	0	150	0	0	150	0
Water + SLS	138	12	92.0	146	4	97.3	132	18	88.0
Water Only	135	15	90	144	6	96.0	130	20	86.7

[0042] All samples tested totally suppressed the germination of spores of the three tested fungi with the exception of Sample 4 on *Penicillium* sp.

[0043] The invention generally describes a synergistic effect when garlic oil or extract is combined with essential oils which results in an improved insecticide/fungicide which is natural and contains no chemical additives. Essential oils are defined in this application to be volatile liquids obtained from plants and seeds including cotton seed oil, soybean oil, cinnamon oil, corn oil, cedar oil, castor oil, clove oil, geranium oil, lemongrass oil, linseed oil, mint oil, sesame oil, thyme oil, rosemary oil, anise oil, basil oil, camphor oil, citronella oil, eucalyptus oil, fennel oil, ginger oil, grapefruit oil, lemon oil, mandarin oil, orange oil, pine needle oil, pepper oil, rose oil, tangerine oil, tea tree oil, tee seed oil, mineral oil and fish oil.

[0044] The active ingredients of garlic oil or garlic extract may be produced and used in the invention, e.g. by manufacture or separated from garlic.

Claims

1. A composition of a pesticide, insecticide and/or fungicide, comprising component (a) garlic oil or garlic extract; and component (b) a second oil selected from the group consisting of essential oils, mineral oils, fish oils and mixtures thereof;
wherein the components (a) and (b) are contained in the composition in a synergistic amount such that the composition is more effective at controlling the growth of bacteria, fungi or insects than the equivalent amounts of components (a) and (b) when used separately.
2. The composition according to claim 1, where said essential oil is selected from the group consisting of cotton seed oil, soybean oil, cinnamon oil, corn oil, cedar oil, castor oil, clove oil, geranium oil, lemongrass oil, linseed oil, mint oil, sesame oil, thyme oil, rosemary oil, anise oil, basil oil, camphor oil, citronella oil, eucalyptus oil, fennel oil, ginger oil, grapefruit oil, lemon oil, mandarin oil, orange oil, pine needle oil, pepper oil, rose oil, tangerine oil, tea tree oil and tee seed oil.
3. The composition according to claim 1 or claim 2, wherein component (b) is selected from the group consisting of cotton seed oil, cinnamon oil, and mineral oil.
4. The composition according to any one of claims 1 to 3 wherein component (b) further includes oils selected from the group consisting of soybean oil, corn oil, cedar oil, castor oil, clove oil, geranium oil, lemongrass oil, linseed oil, mint oil, sesame oil, thyme oil, rosemary oil, anise oil, basil oil, camphor oil, citronella oil, eucalyptus oil, fennel oil, ginger oil, grapefruit oil, lemon oil, mandarin oil, orange oil, pine needle oil, pepper oil, rose oil, tangerine oil, tea tree oil, tee seed oil and fish oil.
5. The composition according to any one of claims 1 to 4, wherein the weight ratio of component (a) to component (b) is between 5% to 98% component (a) to 95% to 2% component (b); preferably the weight ratio of component (a) to component (b) is between 10 to 70% component (a) to 90 to 30% component (b) - and more preferably at least 50% by weight of the total weight of components (a) and (b) is component (a) e.g. 60%.
6. The composition according to any one of the preceding claims wherein either the composition comprises:
a fungicide and the composition is more effective for inhibiting the growth of *Rhizoctonia* sp., *Trichoderma* sp., and *Botrytis* sp. than if an equivalent amount of either component (a) or said (b) were used alone; or a pesticide and the composition is more effective for inhibiting the growth of powdery mildew disease on plants than if an equivalent amount of either component (a) or (b) were used alone.
7. The composition according to any one of the preceding claim, wherein the composition is mixed with water before use and preferably the composition is more effective when compared to the individual components when mixed with water and optionally wherein water is used in excess, and preferably upto 99 parts by weight of water are used to one part by weight of components (a) and (b), more preferably upto 99 parts by weight of water to one part by weight of components (a) and (b).
8. The composition according to any one of the preceding claims further including an emulsifier to emulsify component (a) in component (b) or vice versa; optionally the emulsifier comprising sodium lauryl sulphate and further optionally the emulsifier being used in upto 10 parts by weight to 90 parts of components (a) and (b).

9. The composition according to any one of the preceding claims; wherein component (a) is garlic extract; and/or component (a) is produced from a natural source.
- 5 10. Use of composition as a pesticide, an insecticide and/or a fungicide, the composition comprising component (a) being garlic oil or garlic extract and component (b) being a second oil selected from the group consisting of essential oils, mineral oils, fish oils, and mixtures thereof, in synergistic amounts and optionally being the composition according to any one of claims 1 to 9.

10

15

20

25

30

35

40

45

50

55



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 99 30 2286

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
X	"NATURAL GARLIC SPRAY -- For Insect Control" 'ONLINE', 16 November 1997, pages 1-4, XP002107679 http://www.cyberstreet.com/garlic/gar_bar.htm Retrieved from the Internet on 1999-06-29 * page 2, paragraph 3 * * page 3, paragraph 4 * ---	1,5-10	A01N65/00 //(A01N65/00, 65:00,63:00, 27:00)
X	DATABASE CABA STN-International STN-accession no. 87:19319, V.FRANTZ: "Insect pest control in the greenhouse: alternatives to commercial toxins" XP002107681 * abstract * & PROCEEDINGS OF THE INDIANA ACADEMY OF SCIENCE, vol. 94, 1984, page 98 ---	1,3,5, 7-10	
			TECHNICAL FIELDS SEARCHED (Int.Cl.6)
X	DATABASE CROPU STN-International STN-accession no. 96-83026, N.MOTOYAMA ET AL.: "Insecticidally active component of the so-called natural plant-extract formulation 'Muso'; used as organic agricultural materials" XP002107682 * abstract * & J.PESTIC.SCI., vol. 21, no. 1, 1996, pages 73-79, --- -/--	1,2,4,5, 7,9,10	A01N
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 29 June 1999	Examiner Lamers, W
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

EPO FORM 1503 03/82 (P04C01)



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 99 30 2286

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
X	BIOLOGICAL ABSTRACTS, vol. 1996, Philadelphia, PA, US; abstract no. 58665, H.M.FLINT ET AL.: "Test of garlic oil for the control of the silverleaf whitefly, Bemisia argentifolii bellows and perring (Homoptera: Aleyrodidae) in cotton" XP002107680 * abstract * & SOUTHWESTERN ENTOMOLOGIST, vol. 20, no. 2, 1995, pages 137-150, ---	1,2,4,5, 7,9,10	TECHNICAL FIELDS SEARCHED (Int.Cl.6)
X	DATABASE EPODOC European Patent Office XP002107683 * abstract * & CN 1 170 504 A (LIU FENG) 21 January 1998 ---	1,3,5-10	
X	DATABASE CROPU STN-International STN-accession no. 97-89041, D.L.KERNS ET AL.: "Control of lepidopterous larvae on lettuce with garlic, 1996" XP002107684 * abstract * & ARTHROPOD.MANAGE.TESTS, vol. 22, 1997, page 135 ---	1,4,5, 7-10	
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 29 June 1999	Examiner Lamers, W
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			

EPO FORM 1503 03/82 (204/C01)



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 99 30 2286

DOCUMENTS CONSIDERED TO BE RELEVANT				
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)	
X	DATABASE CABA STN-International STN-accession no. 97:75697, T.H.BRIGGS ET AL.: "Organic insecticides not as effective as synthetics, but still useful for vegetable insect control" XP002107685 * abstract * & HIGHLIGHTS OF AGRICULTURAL RESEARCH - ALABAMA AGRICULTURAL EXPERIMENT STATION, vol. 43, no. 3, 1996, pages 16-18, ---	1,4,5, 7-10		
X	DATABASE WPI Section Ch, Week 9747 Derwent Publications Ltd., London, GB; Class D13, AN 97-503954 XP002107687 & CN 1 123 624 A (SANHE IND CO LTD QIYANG COUNTY HUNAN PRO), 5 June 1996 * abstract * ---	1,2,4-7, 9,10		
X	DATABASE WPI Section Ch, Week 9750 Derwent Publications Ltd., London, GB; Class D13, AN 97-536790 XP002107688 & CN 1 128 107 A (ZHANG Y), 7 August 1996 * abstract * ---	1,2,4-7, 9,10		TECHNICAL FIELDS SEARCHED (Int.Cl.6)
X	GB 1 074 751 A (CLINICAL BIO-ELECTROMETER) 5 July 1967 * the whole document * --- -/--	1,2,4-7, 9,10		
The present search report has been drawn up for all claims				
Place of search THE HAGUE		Date of completion of the search 29 June 1999	Examiner Lamers, W	
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document				

EPO FORM 1503 03.82 (P04C01)



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 99 30 2286

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
X	DATABASE WPI Section Ch, Week 9744 Derwent Publications Ltd., London, GB; Class D13, AN 97-471649 XP002107689 & CN 1 119 498 A (SANHE IND CO LTD QIYANG COUNTY HUNAN PRO), 3 April 1996 * abstract *	1-7, 9, 10	
X	CA 2 107 649 A (FOTOPOULOS THEMIS) 5 April 1995 * page 3, paragraph 4 * * page 5, paragraph 5 *	1, 3, 5-7, 9, 10	
X	DATABASE WPI Section Ch, Week 9720 Derwent Publications Ltd., London, GB; Class C05, AN 97-221661 XP002107690 & JP 09 067220 A (NIPPON SEIYAKU KK) , 11 March 1997 * abstract *	1, 2, 4-7, 9, 10	
X	DATABASE WPI Section Ch, Week 9717 Derwent Publications Ltd., London, GB; Class B04, AN 97-180065 XP002107691 & CN 1 092 995 A (MI C), 5 October 1994 * abstract *	1, 2, 4-7, 9, 10	
			TECHNICAL FIELDS SEARCHED (Int.Cl.6)
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 29 June 1999	Examiner Lamers, W
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

EPO FORM 1503 (3/12) (PACU1)



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 99 30 2286

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
X	DATABASE CROPU STN-International STN-accession no. 98-85795, R.MONTES BELMONT ET AL.: "Control of Aspergillus flavus in maize with plant essential oils and their components" XP002107686 * abstract * & J.FOOD PROT., vol. 61, no. 5, 1998, pages 616-619, -----	1-7,9,10	
			TECHNICAL FIELDS SEARCHED (Int.Cl.6)
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 29 June 1999	Examiner Lamers, W
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

EPO FORM 1503 03 92 (P/M/C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 99 30 2286

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

29-06-1999

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB 1074751 A		NONE	
CA 2107649 A	05-04-1995	NONE	

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82